

**Midterm Exam, Fall 2006**

## ECSE-6968 – Physical Foundations of Solid-State Devices

- Note:** (i) Put your name on paper, show your work, underline results, and always show units.  
(ii) Textbook, manuscript, excerpts, and calculators are allowed.

- Assume that an electron has a kinetic energy  $E = 0.025 \text{ eV} = 0.025 \times 1.602 \times 10^{-19} \text{ CV}$  and is propagating along the positive  $x$  direction in a constant potential given by  $U(x) = U_0 = 0$ .
  - Calculate the velocity of the electron (give numerical value).
  - Calculate the momentum of the electron (give numerical value).
  - Calculate the de Broglie wavelength of the electron (give numerical value).
  - Assume that the wave function of the electron is given by  $\Psi(x, t) = \Psi_0 \exp(kx - \omega t)$ . Calculate the wave number  $k$  of the wave function (give numerical value).
  - Calculate the temporal frequency ( $\omega$ ) of the wave function (give numerical value).
- Assume that the dispersion relation of an electron in a semiconductor near the band minimum is given by  $E = (9.04 \times 10^{-38} \text{ kg m}^4 \text{ s}^{-2}) k^2$ .
  - Give the value of the curvature of the dispersion relation (give numerical value).
  - Calculate the value effective mass (in kg) of an electron obeying this dispersion relation (give numerical value).

Assume that the dispersion relation of a free electron is given by  $E = 100 \text{ meV} = \hbar^2 k^2 / (2m)$  where  $m = 9.1 \times 10^{-31} \text{ kg}$ .

  - Calculate the particle velocity (give numerical value).
  - Calculate the wave number  $k$  and the wavelength  $\lambda$  of the electron (give numerical value).
  - Assume that the energy  $E$  in the equation  $E = \hbar^2 k^2 / (2m)$  is purely kinetic, so that  $E = \hbar\omega$ . This is valid for free particles. Eliminate  $E$  from the last two equations and solve the resulting equation for  $\omega$ . Calculate the group velocity of the electron with energy  $E = 100 \text{ meV}$  using the equation  $v_{\text{gr}} = d\omega / dk$  (give numerical value).
- Consider an electron in a one-dimensional periodic structure along the  $x$  direction with lattice constant (period)  $a = 5 \text{ \AA} = 0.5 \text{ nm}$ . Assume that there will be two allowed bands with a band width  $2\Delta E_0$  and  $2\Delta E_1$ . Assume further that the dispersion relation of the lowest band is given by  $E = E_0 - \Delta E_0 \cos ka$  where  $E_0 = 20 \text{ meV}$  and  $\Delta E_0 = 10 \text{ meV}$ .
  - Which of the following relations will be correct: (i)  $\Delta E_0 > \Delta E_1$  (ii)  $\Delta E_0 < \Delta E_1$  (iii) cannot be determined.
  - What is the effective mass of the electron in the lowest band at the Brillouin center (give numerical value).
  - Let us assume that the electron in the periodic lattice potential is accelerated by an electric field  $\mathcal{E} = 1000 \text{ V/cm}$  along the positive  $x$  direction. After sufficiently long time, the mass of the electron becomes heavier. Give a reason for the fact that the electron mass increases.
  - Give an estimate of the time it takes for the electron effective mass to become infinitely large.
- An electron with effective mass  $m^* = 0.067 m_0$  tunnels through a barrier that is  $100 \text{ meV}$  high and  $L_B = 100 \text{ \AA} = 10 \text{ nm}$  wide.
  - Give a formula that allows one to calculate the tunneling probability of the electron.
  - Calculate the tunneling probability of the electron (give numerical value).
  - Consider that the effective mass of the electron doubles. Does the tunneling probability (i) increase or (ii) decrease?
  - Calculate the tunneling probability of the twice-as-heavy electron (give numerical value).